

*Oïdes maculata*, Oliv.*Oïdes subhemisphærica*, Guér.*Oïdes indica*, Baly.

Judging from the description and figure given by Olivier and from the description given by Guérin, the two species indicated above are, I think, identical with Baly's *Oïdes indica*, the type of which is before me. I am unable to find any definite characters by which to separate specimens from Java from those coming from other localities, such as Tringano, Siam, Burmah, and Assam.

LI.—*Descriptions of two new Species of Sandwich-Island Birds.* By SCOTT B. WILSON, F.Z.S.

*Himatione mana*, sp. n.

*Male.* Head ashy olive, shading into dull olive-green on the back; beneath dull greenish buff, except the chin and throat, which are whitish; wings and tail brown, edged outwardly with olive-green.

*Female.* Duller on the upper parts, while beneath the chin and throat are nearly white, the rest of the underparts more buff than in the male.

*Dimensions.* Total length 4.45 inches, wing 2.50, culmen .45, tarsus .70, tail 1.40.

*Hab.* Hawaii.

*Obs.* The bill is nearly straight, being almost similar to that of *Oreomyza*, while the plumage of the underparts, especially of the female, bears some resemblance to that species. It has not, however, the rounded wing and very short tail of *Oreomyza*.

*Pheornis lanaiensis*, sp. n.

This species closely resembles *Ph. obscura* and *Ph. myiastina*, but is smaller in dimensions than either; while the bill is distinctly intermediate in size between those two species. The outer pair of tail-feathers only have very slight white markings at the tip, but the abdomen and under tail-coverts are nearly pure white.

The length of the wing from the carpal joint is only 3.65, as against 4 of *Ph. obscura*.

*Hab.* Lanai.

*Obs.* The species found on Molokai seems to be identical with the bird from Lanai here described.